

Green's Function for a Neutron in Interaction with a Straight Current Carrying Wire*

Rafael de Lima Rodrigues^{a)}

Departamento de Ciências Exatas e da Natureza

Universidade Federal da Paraíba

58.900-000 – Cajazeiras, PB – Brazil

Departamento de Física Universidade Federal da Paraíba

58.100-000 – Campina Grande, PB – Brazil

and

Arvind Narayan Vaidya^{b)}

Instituto de Física

Universidade Federal do Rio de Janeiro

21.945-970 - Rio de Janeiro, RJ – Brazil

Abstract

We construct an integral representation for the momentum space Green's function for a Neutron in interaction with a straight current carrying wire.

PACS numbers: 11.30.Pb, 03.65.Fd, 11.10.Ef

*This research was supported in part by the Brazilian Research Agencies CNPq and Finep.

Consider an electrically neutral spin 1/2 particle of mass M and a magnetic moment $\mu\vec{\sigma}$ (a Neutron) in interaction with an infinite straight wire carrying a current I and located along the z -axis. The magnetic field generated by the wire is given by (we use units with $c = 1 = \hbar$)

$$\vec{B} = 2I \frac{(-y, x, 0)}{(x^2 + y^2)^{3/2}} \quad (1)$$

where x, y are cartesian coordinates in the plane perpendicular to the wire. The Hamiltonian of the particle is given by

$$H = \frac{\vec{p}^2}{2m} + \mu\vec{\sigma} \cdot \vec{B}. \quad (2)$$

The motion along the z -axis is free and will be ignored in the following. Thus we get a two-dimensional problem with

$$H = \frac{\vec{p}^2}{2m} + 2I\mu \frac{(-y\sigma_1 + x\sigma_2)}{(x^2 + y^2)^{3/2}}. \quad (3)$$

The problem formulated above has a dynamical symmetry [1]. From the form of the Hamiltonian there is one obvious constant of motion

$$J_3 = xp_y - yp_x + \frac{\sigma_3}{2}. \quad (4)$$

Also, the ‘Runge-Lenz vector’ defined by

$$A_i = \frac{\varepsilon_{ij}x_j}{\sigma_1x_2 - \sigma_2x_1} + \frac{1}{4I\mu M}[p_i, J_3]_+, \quad (i = 1, 2) \quad (5)$$

satisfies the following commutation rules

$$[J_3, A_i] = i \varepsilon_{ij} A_j, \quad [A_i, H] = 0, \quad [A_i, A_j] = -i H J_3 \frac{2M}{(2I\mu M)^2}. \quad (6)$$

Thus for negative energy ($-E$) eigenstates of H we may put

$$J_i = A_i \left\{ \frac{(2I\mu M)^2}{-2M E} \right\}^{1/2}, \quad (i = 1, 2) \quad (7)$$

we get

$$[J_\alpha, J_\beta] = i \varepsilon_{\alpha\beta\gamma} J_\gamma \quad (8)$$

where $\alpha, \beta = 1, 2, 3$. Thus J_α generate the algebra of $SO(3)$ in the space of negative energy eigenstates of H . For the space of positive energy eigenstates of H we change $-E$ to $+E$ in above and get the commutation rules of the $SO(2,1)$ algebra.

The $SO(3)$ dynamical symmetry leads to an accidental degeneracy in the bound state spectrum. One can show that ($E < 0$)

$$\vec{A}^2 = 1 + \frac{2M}{(2I\mu M)^2} \left(J_3^2 + \frac{1}{4} \right) \quad (9)$$

and

$$H = -\frac{(2I\mu M)^2}{2M(\vec{J}^2 + \frac{1}{4})} \quad (10)$$

so that the bound state spectrum is given by

$$E_{j+\frac{1}{2}, M} = -\frac{(2I\mu M)^2}{2M(j + \frac{1}{2})^2} \quad (11)$$

where $j = \frac{1}{2}, \frac{3}{2}, \dots$ and $m = -j, -j+1, \dots, j$; which exhibits the $(2j+1)$ fold degeneracy explicitly.

It may be mentioned that even if the bound state spectrum is deduced on the basis of dynamical symmetry it is difficult to use it for the construction of wave functions in the coordinate representation which have an awkward form. As in the case of the Coulomb problem the momentum representation allows us to use the dynamical symmetry in an elegant manner.

As an example we formulate and solve the integral equation for the Green's function $G(\vec{p}, \vec{p}')$ in momentum space. The calculation follows Schwinger's [4] technique for the Coulomb Green's function except for one important modification.

To go over to the momentum representation we put

$$\vec{r} = i \vec{\nabla}_p \quad (12)$$

and use

$$\nabla_p^2 \frac{1}{2\pi} \ln |\vec{p} - \vec{p}'| = \delta(\vec{p} - \vec{p}') \quad (13)$$

so that

$$\frac{i\vec{\nabla}_p}{\nabla_p^2} \Phi(\vec{p}) = \frac{i}{2\pi} \int \Phi(\vec{p}') \frac{(\vec{p} - \vec{p}')}{|\vec{p} - \vec{p}'|^2} d^2 p'. \quad (14)$$

The momentum space Green's function $G(\vec{p}, \vec{p}')$ for energy E satisfies

$$(E - \frac{\vec{p}^2}{2M})G(\vec{p}, \vec{p}') - \frac{iI\mu}{\pi} \int d^2 p'' \frac{\sigma_i \varepsilon_{ij}(\vec{p} - \vec{p}'')_j}{|\vec{p} - \vec{p}''|^2} G(\vec{p}'', \vec{p}') = \delta(\vec{p} - \vec{p}') \quad (15)$$

where $\sigma_i (i = 1, 2, 3)$ are the well known Pauli matrices and $\varepsilon_{12} = 1 = -\varepsilon_{21}, \varepsilon_{ii} = 0$. Assuming that

$$E = -\frac{p_0^2}{2M} \quad (16)$$

is real and negative ($p_0 > 0$) we use the coordinates (n_0, n_i) where

$$n_0 = \frac{p_0^2 - \vec{p}^2}{p_0^2 + \vec{p}^2}, \quad n_i = \frac{2p_0 p_i}{p_0^2 + \vec{p}^2} \quad (17)$$

and

$$n_0^2 + \vec{n}^2 = 1. \quad (18)$$

It is convenient to treat $(n_0, \vec{n})$ as a three dimensional vector n . Here and in the following three dimensional vectors will be written without the vector symbol. The scalar product of two 3-vectors a and b will be written as $a.b$, where $a = (a_0, a_1, a_2)$.

The element of area (solid angle) on the surface the unit sphere of equation (15) is

$$d\Omega = \frac{dn_1 dn_2}{n_0} \Big|_{|n|=1}. \quad (19)$$

Now

$$dn_1 dn_2 = d^2 p \quad J \left(\frac{p_i}{n_j} \right) \quad (20)$$

where the Jacobian $J \left(\frac{p_i}{n_j} \right)$ is given by

$$J\left(\frac{p_i}{n_j}\right) = \left(\frac{2p_0}{p_0^2 + \vec{p}^2}\right)^2 \det(\mathbf{1} - M) \quad (21)$$

where the matrix elements of M are

$$M_{ij} = \frac{2p_i p_j}{p_0^2 + \vec{p}^2}. \quad (22)$$

Since for a matrix N

$$\ln \det N = \text{tr} \ln N \quad (23)$$

we get

$$\ln \det(\mathbf{1} - M) = \ln n_0 \quad (24)$$

where we use the property

$$M^2 = M(1 - n_0). \quad (25)$$

Thus

$$d\Omega = \left(\frac{2p_0}{p_0^2 + \vec{p}^2}\right)^2 d^2p \quad (26)$$

which gives

$$\delta(\Omega - \Omega') = \left(\frac{p_0^2 + \vec{p}^2}{2p_0}\right)^2 \delta(\vec{p} - \vec{p}'). \quad (27)$$

Further

$$\begin{aligned} (n - n')^2 &= (n_0 - n'_0)^2 + (\vec{n} - \vec{n}')^2 \\ &= \frac{(1 + n_0)(1 + n'_0)}{p_0^2} (\vec{p} - \vec{p}')^2. \end{aligned} \quad (28)$$

We define

$$\Gamma(\Omega, \Omega') = -\frac{p_0^4}{M(1 + n_0)^{3/2}(1 + n'_0)^{3/2}} \hat{U}^{-1}(n) G(\vec{p}, \vec{p}') \hat{U}(n') \quad (29)$$

and

$$D(\Omega, \Omega') = -\frac{1}{4\pi} \hat{U}^{-1}(n) \frac{2i(\vec{\sigma} \wedge (\vec{p} - \vec{p}'))}{p_0} \frac{(1+n_0)^{1/2}(1+n'_0)^{\frac{1}{2}}}{(n-n')^2} \hat{U}(n') \quad (30)$$

We get

$$\Gamma(\Omega, \Omega') - \nu \int D(\Omega, \Omega'') \Gamma(\Omega'', \Omega') d\Omega'' = \delta(\Omega - \Omega') \quad (31)$$

where

$$\nu = \frac{2I\mu M}{p_0} \quad (32)$$

which *looks* covariant under 3-dimensional rotations due to the notational artifice.

The introduction of the matrices $\hat{U}(n)$ although permitted would appear to be redundant. However, if we omit them the form of $D(\Omega, \Omega')$ would not display in a convenient fashion the dependence on the 3-vectors n and n' , although the left hand side of equation (33) should do so. We determine $\hat{U}(n)$ by having $D(\Omega, \Omega')$ satisfy a differential equation where its rotational invariance is obvious.

We note that

$$\frac{\vec{\sigma} \wedge (\vec{p} - \vec{p}')}{(n - n')^2 p_0^2} (1 + n_0)(1 + n'_0) = \frac{\vec{\sigma} \wedge (\vec{p} - \vec{p}')}{|\vec{p} - \vec{p}'|^2}. \quad (33)$$

Also

$$\begin{aligned} (\vec{\sigma} \wedge \vec{\nabla}_p) \frac{(\vec{\sigma} \wedge (\vec{p} - \vec{p}'))}{|\vec{p} - \vec{p}'|^2} &= 2\pi \delta(\vec{p} - \vec{p}') \\ &= 2\pi \left(\frac{(1 + n_0)}{p_0} \right)^2 \delta(\Omega - \Omega'). \end{aligned} \quad (34)$$

Thus

$$(\vec{\sigma} \wedge \vec{\nabla}_p) (1 + n_0)^{\frac{1}{2}} (1 + n'_0)^{\frac{1}{2}} \hat{U}(n) D \hat{U}^{-1}(n') = -ip_0 \left(\frac{1 + n_0}{p_0} \right)^2 \delta(\Omega - \Omega'). \quad (35)$$

Let

$$\hat{U}(n) = \sqrt{\frac{1 + n_0}{2}} \hat{U}(n) \quad (36)$$

then

$$\hat{U}^{-1}(n) \frac{ip_0}{1+n_0} (\vec{\sigma} \wedge \vec{\nabla}_p) \hat{U}(n) D(\Omega, \Omega') = \delta(\Omega - \Omega'). \quad (37)$$

The right hand side of equation (37) suggests the use of polar coordinates, hence if the polar angles of $(n_0, \vec{n})$ are θ, ϕ ;

$$\begin{aligned} p_1 &= p_0 \tan \left(\frac{\theta}{2} \right) \cos \phi, \\ p_2 &= p_0 \tan \left(\frac{\theta}{2} \right) \sin \phi. \end{aligned} \quad (38)$$

We then have

$$\frac{ip_0}{1+n_0} \vec{\sigma} \wedge \vec{\nabla}_p = -i(\sigma \cdot e_\phi \partial_\theta - \frac{\sigma \cdot e_p}{\sin \theta} \partial_\phi) \quad (39)$$

where e_p and e_ϕ are unit vectors.

Thus the operator acting on $D(\Omega, \Omega')$ in equation (39) involves at most first order differential operators ∂_θ and ∂_ϕ . If it were invariant under rotations we can make $D(\Omega, \Omega')$ invariant too. The unique candidate in that case is the operator $a + b\sigma \cdot L$ where a and b are constants. We choose $\hat{U}(n)$ so that this holds true.

Since

$$\sigma \cdot L = -i(\sigma \cdot e_\phi \partial_\theta + \sigma_3 \partial_\phi - \cot(\theta) + \sigma \cdot e_p \partial_\phi) \quad (40)$$

the choice

$$\hat{U}(n) = f(\theta, \phi) \exp \left(\frac{ig(\theta, \phi)}{2} \sigma \cdot e_\phi \right) \quad (41)$$

maintains the $-i\sigma \cdot e_\phi \partial_\theta$ term (apart from other additional terms without derivatives).

Working out the derivatives it is easy to see that one must choose $g(\theta, \phi) = \theta$ which gives

$$\begin{aligned} &\hat{U}^{-1} \left(-i\vec{\sigma} \cdot \vec{e}_\phi \partial_\phi + i \frac{\vec{\sigma} \cdot \vec{e}_p}{\sin \theta} \partial_\phi \right) \hat{U} \\ &= \sigma \cdot L - i\hat{U}^{-1} \sigma \cdot e_\phi (\partial_\theta \hat{U}) + i \hat{U}^{-1} \frac{\sigma \cdot e_p}{\sin \theta} (\partial_\phi \hat{U}) \end{aligned} \quad (42)$$

where in the bracketed terms the derivatives act only on $\hat{U}$.

If we impose the condition that the last two terms are independent of the σ -matrices we get

$$\hat{U}(n) = \frac{1+n_0}{2} + \frac{i}{2} \sin\theta \sigma.e_\phi \quad (43)$$

and then

$$\hat{U}^{-1}(n) \frac{ip_0}{1+n_0} \vec{\sigma} \wedge \vec{\nabla}_p \hat{U}(n) = \mathbf{1} + \sigma.L. \quad (44)$$

Hence

$$(\mathbf{1} + \sigma.L)D(\Omega, \Omega') = \delta(\Omega, \Omega'). \quad (45)$$

Our choice of $\hat{U}(n)$ gives

$$\hat{U}(n) = \sqrt{\frac{1+n_0}{2}} \left(1 + \frac{\sigma_3 \vec{\sigma} \cdot \vec{p}}{p_0} \right) \quad (46)$$

which allows us to calculate $D(\Omega, \Omega')$ explicitly by using equation (32). We get

$$D(\Omega, \Omega') = \frac{1}{2\pi} \frac{(1 - \sigma.n \sigma.n')}{(n - n')^2}. \quad (47)$$

Although we know $D(\Omega, \Omega')$ in a closed form it is convenient to rewrite it in terms of the eigenfunctions of the operator $\mathbf{1} + \sigma.L$. The eigenfunctions of the operators J^2, L^2, S^2 and J_3 where $S = \frac{\sigma}{2}$ and $J = L + S$ are known to be $Y_{jm}^\pm(\theta, \phi)$ where $\pm$ correspond to $j = \ell \pm \frac{1}{2}$ respectively [5]. Also

$$(\mathbf{1} + \sigma.L)Y_{jm}^\pm(\theta, \phi) = \pm(j + \frac{1}{2})Y_{jm}^\pm(\theta, \phi) \quad (48)$$

so that

$$D(\Omega, \Omega') = \sum_{jm} \frac{1}{j + \frac{1}{2}} \left\{ Y_{jm}^+(\theta, \phi) Y_{jm}^{+\dagger}(\theta', \phi') - Y_{jm}^-(\theta, \phi) Y_{jm}^{-\dagger}(\theta', \phi') \right\}. \quad (49)$$

Going back to equation (33) we get

$$\Gamma(\Omega, \Omega') = \sum_{jm} \left\{ \frac{Y_{jm}^+(\theta, \phi) Y_{jm}^{+\dagger}(\theta', \phi')}{1 - \frac{\nu}{j + \frac{1}{2}}} + \frac{Y_{jm}^-(\theta, \phi) Y_{jm}^{-\dagger}(\theta', \phi')}{1 + \frac{\nu}{j + \frac{1}{2}}} \right\} \quad (50)$$

which for $\nu > 0$ has poles at

$$\nu = \frac{2IM\mu}{p_0} = j + \frac{1}{2}. \quad (51)$$

This gives the bound state energies which are same as before. The unnormalized bound state wave functions are $\frac{1}{(1+n_0)^{3/2}} \hat{U}^{-1} Y_{jm}^+(\theta, \phi)$.

Next, one can do the sum indicated in equation (52) to write an integral representation for $\Gamma(\Omega, \Omega')$ which in turn leads to one for the Green's function $G(\vec{p}, \vec{p}')$.

This is done by using the identities:

$$\frac{1}{1 \mp \frac{\nu}{j+\frac{1}{2}}} = \frac{j+\frac{1}{2}}{j+\frac{1}{2} \mp \nu} = 1 \pm \frac{\nu}{j+\frac{1}{2} \mp \nu} = 1 \pm \frac{\nu}{j+\frac{1}{2}} + \frac{\nu^2}{(j+\frac{1}{2})(j+\frac{1}{2} \mp \nu)} \quad (52)$$

and using

$$\frac{1}{n} = \int_0^1 \rho^{n-1} d\rho \quad (53)$$

valid for $n > 1$. Although each form of the identity leads to a different looking integral representation they are related through integration by parts. We concentrate on the second form.

We have

$$\Gamma(\Omega, \Omega') = \delta(\Omega, \Omega') + \nu \int_0^1 (\rho^{-\nu} \Delta^+ - \rho^\nu \Delta^-) d\rho \quad (54)$$

where

$$\Delta^\pm = \sum_{jm} \rho^{j-\frac{1}{2}} Y_{jm}^\pm(\theta, \phi) Y_{jm}^{\pm\dagger}(\theta', \phi'). \quad (55)$$

Using the explicit forms of $Y_{jm}^\pm(\theta, \phi)$ one can show that

$$\begin{aligned} \Delta^+ &= \frac{1}{4\pi} (\partial_\rho \rho + \sigma.L) \Phi, \\ \Delta^- &= \frac{1}{4\pi} \left(-\frac{1}{\rho} \sigma.L + \partial_\rho \right) \Phi \end{aligned} \quad (56)$$

where

$$\Phi = \left((1-\rho)^2 + \rho(n-n')^2 \right)^{-\frac{1}{2}}. \quad (57)$$

Calculating the derivatives we get

$$\begin{aligned}\Delta^+ &= \frac{1}{4\pi}(1 - \rho \sigma.n \sigma.n')\Phi^3, \\ \Delta^- &= \frac{1}{4\pi}(\rho - \sigma.n \sigma.n')\Phi^3.\end{aligned}\tag{58}$$

Thus

$$\Gamma(\Omega, \Omega') = \delta(\Omega, \Omega') + \frac{\nu}{4\pi} \int_0^1 d\rho \left(\rho^{-\nu}(1 - \rho \sigma.n \sigma.n') - \rho^\nu(\sigma.n \sigma.n' - \rho) \right).\tag{59}$$

Substituting this in equation (31) we get the integral representation for $G(\vec{p}, \vec{p}')$:

$$\begin{aligned}G(\vec{p}, \vec{p}') &= \frac{\delta(\vec{p} - \vec{p}')}{E - T} + \frac{\nu}{8\pi M} \frac{1}{E - T} \frac{1}{E - T'} \frac{1}{|\vec{p} - \vec{p}'|^2} \\ &\left\{ \left(2M(E - \frac{T + T'}{2}) - i\sigma_3(\vec{p} \wedge \vec{p}') + \frac{1}{2}|\vec{p} - \vec{p}'|^2 \right) \right. \\ &(I(-\nu) - I(\nu) + I(1 + \nu) - I(1 - \nu)) \\ &\left. + ip_0 \vec{\sigma} \wedge (\vec{p} - \vec{p}')(I(\nu) + I(-\nu) + I(1 + \nu) + I(1 - \nu)) \right\}\end{aligned}$$

where p_0 and ν are given by equations (20) and (34). Also,

$$I(\nu) = \int_0^1 d\rho \frac{\rho^\nu (n - n')^2}{\{(1 - \rho)^2 + \rho(n - n')^2\}^{\frac{3}{2}}},\tag{60}$$

$$T = \frac{p^2}{2M}, \quad T' = \frac{p'^2}{2M}\tag{61}$$

and

$$(n - n')^2 = -\frac{2E}{M} \frac{|\vec{p} - \vec{p}'|^2}{(E - T)(E - T')}.\tag{62}$$

In conclusion we have obtained an integral representation for the Green's function for a Neutron in interaction with a linear current. The results were deduced for the case $E < 0$. By analytic continuation to $E > 0$ one can identify the scattering amplitude if the distorted free propagators can be identified. In contrast to the Coulomb problem it is not easy to solve the scattering problem in the coordinate representation by separation of variables. Although a general momentum transfer dependence of $(\vec{p} - \vec{p}')^{-2}$ is indicated many details need to be worked out. This will be done elsewhere.

REFERENCES

- ^{a)} E-mail: CENDFI76@BRUFPB (BITNET)
- ^{b)} E-mail: IFT10034@UFRJ (BITNET)
- [1] G. P. Pron'ko and Yu G. Stroganov, *Sov. Phys. JETP* **45** (1977) 1073.
- [2] R. Blümel and K. Dietrich, *Phys. Rev. A* **43** (1991) 22.
- [3] I. Voronin, *Phys. Rev. A* **43** (1991) 29.
- [4] J. Schwinger, *J. Math. Phys.* **5** (1964) 1607.
- [5] J. D. Bjorken and S. D. Drell, *Relativistic Quantum Mechanics (New York, McGraw Hill, 1965)*.